


Prepare and Consent
the patient

Where possible
request the patient
cleans their
hand and arm

Preparation zone



1
Clean hands
with alcohol hand rub
or soap & water



2
Clean tray
according to local
policy creating a
General Aseptic Field;
whilst it dries . . .



3
Gather equipment



4
Clean hands
with alcohol hand
rub or soap & water



5
Prepare Equipment
protecting Key-Parts with
non touch technique (NTT)
and Micro Critical Aseptic
Fields (Caps & Covers)

Patient Zone



6
Position arm
& apply apron



7
Apply disposable
tourniquet, locate vein,
release tourniquet



8
Clean hands
with alcohol hand
rub or soap & water



9
Re-tighten
tourniquet



10
Apply gloves



11
Clean site for 30 sec
using a 2%chlorhexidine/
70% alcohol applicator &
across hatch technique;
and allow to dry



12
Anchor vein below
puncture site & insert
cannula using NTT
& secure



13
Using NTT, attach
extension set, flush
device, use a sterile semi-
permeable dressing &
a fixation device



14
Dispose of
sharps
and equipment



15
Dispose of gloves
then apron and
immediately.....



16
Clean hands
with alcohol hand rub
or soap & water

Decontamination Zone



17
Clean tray
according to local
policy



18
Clean hands
with alcohol hand
rub or soap & water