



When entering the patient's home disinfect hands



1 Disinfect Tray
as local policy – creating a **General Aseptic Field**



2 Apply Disposable Apron




3 Gather Equipment
& place around tray



4 Disinfect Hands
with alcohol hand rub or soap & water



5 Prepare Equipment
protecting **Key-Parts** with non-touch technique (NTT) & **Micro Critical Aseptic Fields**



6 Place Drape
under the patient's arm



7 Apply Tourniquet
(use single-use disposable tourniquet) palpate vein & release tourniquet



8 Disinfect Hands
with alcohol hand rub or soap & water



9 Apply Gloves
(use sterile gloves if **Key-Site** must be touched directly)* & re-apply tourniquet



10 Disinfect Skin
with a 2% CHG / 70% IPA using friction for 30 seconds. Allow to dry



11 Access Vein
protecting **Key-Parts** & **Key-Sites** using NTT
*If re-palpation needed, repeat step 10 or use sterile gloves**

If attempt to draw blood is unsuccessful return to step **7**



12 Dispose of Sharps
and equipment



13 Disinfect Tray
according to local policy



14 Dispose of Gloves
then apron & immediately...



15 Disinfect Hands
with alcohol hand rub or soap & water



16 Dispose of Waste
according to local policy



When leaving the patient's home disinfect hands

